



My Gratitude
JOURNAL

So good to see the regulars at exercise class again.

Looking forward to my next class.

Actor portrayal.

Getting started with MYQORZO™

FOR ADULTS WITH SYMPTOMATIC OBSTRUCTIVE HYPERTROPHIC CARDIOMYOPATHY (oHCM) TO IMPROVE FUNCTIONAL CAPACITY AND SYMPTOMS

It is not known if MYQORZO is safe and effective in children.

You're about to take the first steps to starting treatment with MYQORZO. You might be wondering what's next. This brochure is designed to help you and your loved ones know what to expect while starting and taking MYQORZO.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **Heart failure**, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death. You must have echocardiograms (echos) before and during treatment with MYQORZO and monitor for signs and symptoms of heart failure. People who develop a serious illness such as a serious infection or who develop a new or worsening irregular heartbeat have a greater risk of heart failure during treatment with MYQORZO

Tell your healthcare provider or get medical help right away if you develop new or worsening shortness of breath, chest pain, fatigue, leg swelling, a racing sensation in your heart (palpitations), or rapid weight gain.

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Medication Guide**.

 **MYQORZO™**
(aficamten) 5-10-15-20 mg
tablets



Actor portrayal. MYQORZO is not approved for use in people younger than 18 years.

GET READY

Before getting your prescription



Once you and your healthcare provider have determined together that MYQORZO™ may be right for you, here are some things you may expect

Some of these steps you may have already taken with your healthcare provider at your last visit, and others may still need to be done.

✔ **Your healthcare provider will talk with you about your medical status, including if you:**

- Are pregnant or plan to become pregnant, as it is not known if MYQORZO can cause harm to your unborn baby
- Are breastfeeding or planning to breastfeed, as it is not known if MYQORZO passes into breast milk
- Are taking any medicines, including prescription and over-the-counter medicines, vitamins, and herbal supplements

✔ **Your healthcare provider will discuss the MYQORZO Risk Evaluation and Mitigation Strategy (REMS) program.** To help protect the health of people taking medicines that have certain serious risks, the U.S. Food and Drug Administration (FDA) sometimes requires a REMS program. In some people, MYQORZO can increase the risk of heart failure. The MYQORZO REMS program is designed to help reinforce safe use by providing special instructions, monitoring, and provider and patient education.

In order for you to start and continue taking MYQORZO, your healthcare provider will:

- **Have you get echocardiograms to make sure you are able to take MYQORZO**
It is important that you complete your echocardiograms at the times recommended by your doctor
- **Help you complete the Patient Enrollment Form** (required for you to start taking MYQORZO)
Your provider will also discuss the risks and benefits of MYQORZO treatment

✔ **Your healthcare team** can walk you through the full treatment process, or you can refer to the overview on pages 10–11.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **The risk of heart failure is also increased when MYQORZO is taken with certain other medicines.**
Tell your healthcare provider about any prescribed and over-the-counter medicines you take, before and during your treatment with MYQORZO

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Medication Guide**.



GET SET

Getting started on MYQORZO™



MYQORZO & You Patient Support Program

The MYQORZO & You Patient Support Program offers personalized patient support for your treatment journey.

Your healthcare provider can enroll you or you can call 833-MYQORZO (833-697-6796) to learn more.

Affordability assistance

MYQORZO & You offers coverage and affordability programs designed to support you throughout your treatment journey.

MYQORZO & You Copay Savings Program

If you are commercially insured and meet eligibility requirements,* you may pay as little as \$5 for MYQORZO and as little as \$0 for your echocardiograms. You can enroll by visiting myqorzo.com/copaysignup.

Specialty pharmacy

MYQORZO is available through specialty pharmacies that will ship your medication directly to your home. The specialty pharmacy will call you to schedule shipment of your prescription—this may appear as an unfamiliar number on your phone. Make sure to answer the call so that you can get your prescription.



To enroll in the MYQORZO & You Copay Savings Program, visit myqorzo.com/copaysignup.

Eligible patients may pay as little as \$5 for MYQORZO and \$0 for echocardiograms.*

*See program terms and conditions at myqorzo.com.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- Because of the risk of heart failure, MYQORZO is only available through a restricted distribution program called the MYQORZO Risk Evaluation and Mitigation Strategy (REMS) Program
 - Your healthcare provider must be enrolled in the MYQORZO REMS Program for you to be prescribed MYQORZO
 - Before you start treatment with MYQORZO, you must enroll in the MYQORZO REMS Program. Talk to your healthcare provider about how to enroll in the program. You will be given information about the program when you enroll
 - Before you take MYQORZO, your healthcare provider and pharmacist will make sure you understand how to take MYQORZO safely, which will include returning for echos when advised by your healthcare provider. MYQORZO can only be dispensed by a certified pharmacy that participates in the MYQORZO REMS Program
 - If you have any questions about the MYQORZO REMS Program, ask your healthcare provider, go to www.MYQORZOREMS.com, or call 1-844-285-7367

Meet your MYQORZO Navigator

When you enroll in the MYQORZO & You Patient Support Program, you will be connected with a dedicated MYQORZO Navigator. They are here to help you start and stay on your MYQORZO treatment.



Once enrolled in MYQORZO & You, your dedicated MYQORZO Navigator will:



Call to welcome you into the program within 1 business day



Explain affordability support options that may best suit your needs



Provide REMS education and support



Help with shipment coordination of MYQORZO from your specialty pharmacy



Provide ongoing refill reminders and lifestyle support while on MYQORZO



If you have not been enrolled in MYQORZO & You and would still like to request MYQORZO Navigator support, visit MYQORZONavigator.com.



For general support and assistance with access and affordability, contact a MYQORZO Navigator at 833-MYQORZO (833-697-6796) Monday through Friday, 8 AM to 8 PM ET.

For medical questions, call your healthcare provider.

IMPORTANT SAFETY INFORMATION (continued)

Who should not take MYQORZO?

Do not take MYQORZO if you take a medicine called rifampin.

What are the possible side effects of MYQORZO?

MYQORZO can cause serious side effects, including heart failure.

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Medication Guide**.






MYQORZO is a cardiac myosin inhibitor (CMI) thoughtfully designed with you in mind

Reach your individualized dose

Your healthcare provider may begin adjusting your dose as early as 2 weeks after starting MYQORZO, reaching your individualized maintenance dose within 8 weeks.*

You have some flexibility in scheduling your echocardiograms (ECHOs). Work with your healthcare provider to schedule an echocardiogram between 2 and 8 weeks after starting MYQORZO and after any dose change.



*Based on echocardiograms performed 2 to 8 weeks after starting treatment and after any dose change.

†MYQORZO may be increased at 5-mg increments up to 20 mg. If at any point you have a drop in your left ventricular ejection fraction (LVEF), experience a severe illness, or start taking certain medications, your healthcare provider may need to lower your dose of MYQORZO or even temporarily stop it.

IMPORTANT SAFETY INFORMATION (continued)

What should I tell my healthcare provider before taking MYQORZO?

Before taking MYQORZO, tell your healthcare provider about all of your medical conditions, including if you:

- Are pregnant or plan to become pregnant. It is not known if MYQORZO can cause harm to your unborn baby. Tell your healthcare provider if you become pregnant during treatment or within 3 weeks after the last dose of MYQORZO. There is a pregnancy study for MYQORZO. Your healthcare provider should report your pregnancy exposure to Cytokinetics, Inc
- Are breastfeeding or plan to breastfeed. It is not known if MYQORZO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with MYQORZO

Once your healthcare provider finds your individualized dose, it is important to take MYQORZO daily, as directed

- Take MYQORZO exactly as your healthcare provider tells you to take it
- Do not change your dose of MYQORZO without talking to your healthcare provider first
- Take MYQORZO 1 time a day, at about the same time each day
- Take MYQORZO with or without meals
- Swallow the MYQORZO tablet whole
- If you miss a dose of MYQORZO, take it as soon as possible and take your next dose at your regularly scheduled time the next day. Do not take 2 doses on the same day to make up for a missed dose
- Your healthcare provider may change your dose, or temporarily stop your treatment with MYQORZO if you have certain side effects
- If you take too much MYQORZO, call your healthcare provider or the Poison Control Help Line at 1-800-222-1222, or go to the nearest hospital emergency room right away

MYQORZO is a once-daily pill and is available in 4 dose strengths:



Pills not actual size.

Size vs a US dime: 5 mg 10 mg 15 mg 20 mg

US dime for scale

IMPORTANT SAFETY INFORMATION

What should I tell my healthcare provider before taking MYQORZO?

Before and during MYQORZO treatment, tell your healthcare provider about all the prescription and over-the-counter medicines, vitamins, and herbal supplements you take. Taking MYQORZO with certain medicines may lead to increased levels of MYQORZO in your blood and increase the risk of heart failure. **Do not stop or change the dose of a medicine or start a new medicine without telling your healthcare provider.**

Especially tell your healthcare provider if you take fluconazole (if used for more than 3 days), voriconazole, or fluvoxamine.

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Medication Guide**.





MYQORZO can cause serious side effects, including heart failure, which is why your doctor will monitor how your heart is functioning.

Tell your healthcare provider or get medical help right away if you develop new or worsening:

- Shortness of breath
- Chest pain
- Swelling in your legs
- Racing sensation in your heart (palpitations)
- Fatigue
- Rapid weight gain



The most common side effect of MYQORZO is high blood pressure (hypertension).

In the clinical trial, no participants experienced treatment interruptions or worsening heart failure events due to a drop in ejection fraction. Ejection fraction compares the amount of blood in your heart chamber to how much blood is pumped out.



Do not take MYQORZO if you take a medicine called rifampin, as it is not safe to take these medicines together

Before and during MYQORZO treatment, tell your healthcare provider about:



All medicines you are taking:

- Including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking MYQORZO with certain medicines may increase your risk of heart failure
- **Do not stop or change the dose of a medicine or start a new medicine without telling your healthcare provider**
- Especially tell your healthcare provider if you take the medicines fluconazole (if used for more than 3 days), voriconazole, or fluvoxamine, as a dose change may be required

All medical conditions, including if you:

- Are pregnant, are planning to become pregnant, or become pregnant within 3 weeks of your last MYQORZO dose. It is not known if MYQORZO can cause harm to your unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if MYQORZO passes into your breast milk

MYQORZO Risk Evaluation and Mitigation Strategy (REMS) Program



To help protect the health of people taking medicines that have certain serious risks, the U.S. Food and Drug Administration (FDA) sometimes requires a REMS program. In some people, MYQORZO can increase the risk of heart failure. The MYQORZO REMS Program is designed to help reinforce safe use by providing special instructions, monitoring, and provider and patient education.

If your healthcare provider determines MYQORZO is right for you, they will help you enroll in the program to get you started. Learn more at [MYQORZO.com/REMS](https://www.myqorzo.com/REMS).

What is a cardiac myosin inhibitor (CMI)?

CMIs work by reducing myosin activity to allow for more normal myosin and actin interactions. This helps prevent the heart from contracting too strongly. CMIs are the only drug class FDA approved for adults with symptomatic obstructive HCM.

Why do I need to have echocardiograms?

An echocardiogram is a type of ultrasound of the heart. It can help your healthcare provider see how well your heart is functioning. This helps your healthcare provider ensure you're taking the best dose for you.

You will need to have an echocardiogram performed before starting MYQORZO (if you haven't had one recently), every time your dose is increased, and then every 6 months after you reach your individualized dose (some people may need to have them every 3 months, but your healthcare provider will let you know if that applies to you).

How is my individualized dose established?

You will start by taking 5 mg of MYQORZO and have an echocardiogram 2 to 8 weeks after. The echocardiogram lets your healthcare provider see the effects MYQORZO is having on your heart. The dose will be increased in 5-mg increments until you're getting the maximum effect with minimal side effects.

What if I accidentally skip a dose?

If you miss a dose of MYQORZO, take it as soon as possible and take your next dose at your regularly scheduled time the next day. Do not take 2 doses on the same day to make up for a missed dose.

What if I have side effects?

Call your healthcare provider if you experience any new or worsening side effects. You can also report side effects to Cytokinetics® at 833-MEDCYTO (1-833-633-2986), or emailing medicalinformation@cytokinetics.com.

IMPORTANT SAFETY INFORMATION (continued)

What are the most common side effects of MYQORZO?

The most common side effect of MYQORZO is high blood pressure (hypertension).

These are not all the possible side effects of MYQORZO. Talk to your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Cytokinetics at 1-833-633-2986.

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Medication Guide**.

What are the potential side effects of treatment with MYQORZO?

MYQORZO may cause heart failure. Heart failure is a condition where the heart cannot pump with enough force. Heart failure is a serious condition that can lead to death. People who develop a serious infection or new or worsening heart rhythm problem may have a greater risk of heart failure during treatment with MYQORZO.

Tell your provider right away if you experience any of the following symptoms:

- Shortness of breath
- Chest pain
- Swelling in your legs
- Racing sensation in your heart (palpitations)
- Fatigue
- Rapid weight gain

The most common side effect of MYQORZO is high blood pressure (hypertension).

Why is REMS required?

REMS, which stands for Risk Evaluation and Mitigation Strategy, is a drug safety program required by the FDA for certain medications. REMS ensures you have the information you need to take your medicine safely.

Is there support available once I'm taking MYQORZO?

Yes. The MYQORZO & You Patient Support Program offers personalized patient support for your treatment journey. You can enroll through your healthcare provider or learn more by calling 833-MYQORZO (833-697-6796).

Is there assistance with the cost of MYQORZO?

Once enrolled in MYQORZO & You, your Navigator may connect you with affordability programs based on your eligibility. Eligible commercially insured patients may also pay as little as \$5 for MYQORZO through the MYQORZO & You Copay Savings Program. You can sign up at [myqorzo.com/copysignup](https://www.myqorzo.com/copysignup).



Getting started and staying on MYQORZO

GET READY

Getting Your Prescription



Your healthcare provider will talk with you about:

- Any medicines you may be taking
- Any existing medical conditions, including pregnancy



Your provider will schedule an echocardiogram to confirm your eligibility for treatment with MYQORZO



Your provider will discuss the Risk Evaluation and Mitigation Strategy (REMS) program with you and:

- Provide you with the MYQORZO Medication Guide and REMS Patient Guide
- Enroll you in the REMS program or refer you to a REMS-certified healthcare professional who can enroll you. To learn more about REMS, visit [MYQORZO.com/REMS](https://myqorzo.com/REMS) or call 844-285-7367



Your healthcare provider can enroll you in the MYQORZO & You Patient Support Program

- MYQORZO & You offers personalized patient support for your treatment journey
- You can call a MYQORZO Navigator to learn more about the program at 833-MYQORZO (833-697-6796)



Once your doctor prescribes MYQORZO:

- If you've been enrolled in the MYQORZO & You Patient Support Program by your provider or care team, you will receive a welcome call from a MYQORZO Navigator within 1 business day. A Navigator is your dedicated contact to help you start and stay on your MYQORZO treatment

Look for a call from a MYQORZO Navigator

- Not enrolled by your healthcare provider or care team? You can request MYQORZO Navigator support by going to [MYQORZONavigator.com](https://myqorzo.com). You may also sign up for the MYQORZO & You Copay Savings Program by going to myqorzo.com/copaysignup



You will receive a call from the specialty pharmacy to coordinate shipment of your MYQORZO

- Make sure to answer the call from your specialty pharmacy. This may come from an unfamiliar phone number

GET SET

Getting Started on MYQORZO

GET GOING

Taking MYQORZO



Reach your individualized dose

Your healthcare provider may begin adjusting your dose as early as 2 weeks after starting MYQORZO, potentially allowing you to reach your individualized maintenance dose within 8 weeks.*

*Based on echocardiograms performed 2 to 8 weeks after starting treatment and after any dose change.



Your care team will schedule a maintenance echocardiogram every 6 months

Depending on results of your echocardiogram, you may need to have follow-up every 3 months.



Continued MYQORZO Navigator support

If enrolled in MYQORZO & You, you will receive periodic check-in calls from your dedicated Navigator about

- Ongoing eligibility in affordability programs
- Refill reminders
- Lifestyle support



Have you done the following?



Get ready

- Talked with your healthcare provider about your MYQORZO and any medicines or medical conditions
- Got an echocardiogram
- Enrolled in the MYQORZO & You Patient Support Program (optional to enroll)



Get set

- Received a call from your MYQORZO Navigator and discussed savings assistance programs you may be eligible for (optional enrollment)
- Received a call from the specialty pharmacy to coordinate shipping of MYQORZO



Get going

- Received your MYQORZO treatment and started your first dose
- Had your healthcare provider schedule your first echocardiogram (and any that follow after dose changes)

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **Heart failure**, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death. You must have echocardiograms (echos) before and during treatment with MYQORZO and monitor for signs and symptoms of heart failure. People who develop a serious illness such as a serious infection or who develop a new or worsening irregular heartbeat have a greater risk of heart failure during treatment with MYQORZO

Tell your healthcare provider or get medical help right away if you develop new or worsening shortness of breath, chest pain, fatigue, leg swelling, a racing sensation in your heart (palpitations), or rapid weight gain.

IMPORTANT SAFETY INFORMATION (continued)


What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **The risk of heart failure is also increased when MYQORZO is taken with certain other medicines.** Tell your healthcare provider about any prescribed and over-the-counter medicines you take, before and during your treatment with MYQORZO

Please see additional Important Safety Information throughout and full Prescribing Information, including Boxed WARNING, and Medication Guide.





**My
Gratitude
JOURNAL**

Being outside with my son feels great. We built a pile of leaves for him to jump in and it made his day!

Looking forward to more time outside together.

Actor portrayal. MYQORZO is not approved for use in people younger than 18 years.

If you have questions or need support getting started on MYQORZO™, we're here to help.

For medical questions, call your healthcare provider. For support and assistance with access and affordability:



Contact a MYQORZO Navigator at 833-MYQORZO (833-697-6796)
Monday through Friday, 8 AM to 8 PM ET.

For more information about the MYQORZO & You Patient Support Program, visit myqorzo.com/support.

INDICATION AND USAGE

MYQORZO is a prescription medicine used to treat adults with symptomatic obstructive cardiomyopathy (oHCM) to improve functional capacity and symptoms.

It is not known if MYQORZO is safe and effective in children.

Please see full [Prescribing Information](#), including [Boxed WARNING](#), and [Medication Guide](#).



CYTOKINETICS® and the CYTOKINETICS C-shaped logo are registered trademarks of Cytokinetics in the U.S. and certain other countries. MYQORZO™, the MYQORZO logo, and the MYQORZO & You logo are trademarks of Cytokinetics in the U.S. © 2026 CYTOKINETICS. All Rights Reserved. US-AFI-00247 01/2026