

STAYING IN STEP WITH MYQORZO™



Every journey forward begins with a step.
As you get familiar with MYQORZO, we're here
to support your progress.

MYQORZO is the latest FDA-approved treatment for adults with symptomatic obstructive hypertrophic cardiomyopathy (oHCM) to improve functional capacity and symptoms.

It is not known if MYQORZO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **Heart failure**, a condition where the heart cannot pump with enough force, is a serious condition that can lead to death. You must have echocardiograms (echos) before and during treatment with MYQORZO and monitor for signs and symptoms of heart failure. People who develop a serious illness such as a serious infection or who develop a new or worsening irregular heartbeat have a greater risk of heart failure during treatment with MYQORZO

Tell your healthcare provider or get medical help right away if you develop new or worsening shortness of breath, chest pain, fatigue, leg swelling, a racing sensation in your heart (palpitations), or rapid weight gain.



Actor portrayal.

Please see full [Prescribing Information](#), including **Boxed WARNING**, and **Medication Guide**.

 **MYQORZO™**
(aficamten) 5-10-15-20 mg
tablets

Get the most from MYQORZO™



Taking MYQORZO as prescribed* helps it work as intended and may support your overall treatment plan. You've got this.



Have you built in your one-pill, **same-time-of-day** habit? ([See tips here!](#))



Missed a dose? Take it as soon as you can that same day, then take your next dose at your normal time the next day. **Do not take two doses of MYQORZO in one day.**



Feeling better? That could mean your medication is doing its job. Keep taking it as prescribed to stay on track. **Not sure you're feeling better yet?** Stay on course—improvements were seen at 12 and 24 weeks for some people on MYQORZO in the clinical trial.



Be sure to **keep up with appointments and echocardiograms**. Hints to help: schedule follow-up appointments before leaving your healthcare provider's office, use digital calendars with alerts, and set reminders 3 days and 1 day before your appointment.



Bonus tip: Make it a habit to record your progress in a journal to help track your results.

*Take MYQORZO with or without meals. Swallow the MYQORZO tablet whole.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **The risk of heart failure is also increased when MYQORZO is taken with certain other medicines.** Tell your healthcare provider about any prescribed and over-the-counter medicines you take, before and during your treatment with MYQORZO

Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.

 **MYQORZO™**
(aficamten) 5-10-15-20 mg
tablets

Have you reached the right dose for you?




MYQORZO™ comes in different strengths, offering you and your healthcare provider options to help manage your oHCM:

- Throughout your treatment journey, it's important to **keep your healthcare provider up to date with how you're doing**, especially if your symptoms are impacting your daily activities
- Conversations with your healthcare provider help ensure your dose is optimized to deliver **symptom improvement and reduced obstruction**
- **Thinking about stopping MYQORZO?** Do not adjust your medication without speaking to your healthcare provider first



If you're still experiencing symptoms, your healthcare provider may need to adjust your dose to ensure you're receiving the maximum symptom relief. To learn more about how to reach the right dose for you, [click here](#).



Check with your healthcare provider before you start, stop, or change medications.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about MYQORZO?

MYQORZO can cause serious side effects, including:

- **Because of the risk of heart failure, MYQORZO is only available through a restricted distribution program called the MYQORZO Risk Evaluation and Mitigation Strategy (REMS) Program**
 - Your healthcare provider must be enrolled in the MYQORZO REMS Program for you to be prescribed MYQORZO
 - Before you start treatment with MYQORZO, you must enroll in the MYQORZO REMS Program. Talk to your healthcare provider about how to enroll in the program. You will be given information about the program when you enroll
 - Before you take MYQORZO, your healthcare provider and pharmacist will make sure you understand how to take MYQORZO safely, which will include returning for echos when advised by your healthcare provider. MYQORZO can only be dispensed by a certified pharmacy that participates in the MYQORZO REMS Program
 - If you have any questions about the MYQORZO REMS Program, ask your healthcare provider, go to www.MYQORZOREMS.com, or call 1-844-285-7367

Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.


 **MYQORZO™**
(aficamten) 5-10-15-20 mg
tablets

Keep in step



Be as consistent with your medication as you are with other parts of your daily care (eg, brushing your teeth, washing your face).

Here are some ideas on how to stick to your routine—even when you're traveling or your schedule changes:



Put your pills in a case with compartments for each day of the week



Pair taking MYQORZO™ with another daily activity, like taking a shower or eating a meal



Use a calendar or medication journal and check off when you take each dose



Leave the medication in a place you'll see it, away from the reach of children

IMPORTANT SAFETY INFORMATION (continued)

Who should not take MYQORZO?

Do not take MYQORZO if you take a medicine called rifampin.

What are the possible side effects of MYQORZO?

MYQORZO can cause serious side effects, including heart failure.

What should I tell my healthcare provider before taking MYQORZO?

Before taking MYQORZO, tell your healthcare provider about all of your medical conditions, including if you:

- Are pregnant or plan to become pregnant. It is not known if MYQORZO can cause harm to your unborn baby. Tell your healthcare provider if you become pregnant during treatment or within 3 weeks after the last dose of MYQORZO. There is a pregnancy study for MYQORZO. Your healthcare provider should report your pregnancy exposure to Cytokinetics, Inc
- Are breastfeeding or plan to breastfeed. It is not known if MYQORZO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with MYQORZO

Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.



Support for your treatment journey



The MYQORZO & You Patient Support Program offers **personalized patient support** for your treatment journey:



Once enrolled in MYQORZO & You, your dedicated MYQORZO™ Navigator will:



Call to welcome you into the program



Explain affordability support options



Help with shipment coordination



Provide lifestyle support



The **copay program** helps eligible commercially insured patients pay as little as \$5 for MYQORZO, and pay as little as \$0 for echocardiograms*

*See program [terms and conditions](#).

IMPORTANT SAFETY INFORMATION (continued)

What should I tell my healthcare provider before taking MYQORZO?

Before and during MYQORZO treatment, tell your healthcare provider about all the prescription and over-the-counter medicines, vitamins, and herbal supplements you take. Taking MYQORZO with certain medicines may lead to increased levels of MYQORZO in your blood and increase the risk of heart failure. **Do not stop or change the dose of a medicine or start a new medicine without telling your healthcare provider.**

Especially tell your healthcare provider if you take fluconazole (if used for more than 3 days), voriconazole, or fluvoxamine.

Please see full [Prescribing Information](#), including **Boxed WARNING**, and [Medication Guide](#).



One more step: join MYQOMMUNITY™



You're not alone. MYQOMMUNITY is here to support you on your journey.



Sign up today to receive helpful information and [stay connected](#) to resources and updates that are tailored to you.

MYQOMMUNITY. Where learning meets inspiration.



Actor portrayal.

IMPORTANT SAFETY INFORMATION (continued)

What are the most common side effects of MYQORZO?

The most common side effect of MYQORZO is high blood pressure (hypertension).

These are not all the possible side effects of MYQORZO. Talk to your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to Cytokinetics at 1-833-633-2986.

Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.





For more information, visit myqorzo.com

INDICATION AND USAGE

MYQORZO is a prescription medicine used to treat adults with symptomatic obstructive hypertrophic cardiomyopathy (oHCM) to improve functional capacity and symptoms.

It is not known if MYQORZO is safe and effective in children.

Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.

CYTOKINETICS® and the CYTOKINETICS C-shaped logo are registered trademarks of Cytokinetics in the U.S. and certain other countries. MYQORZO™, the MYQORZO logo, the MYQORZO & You logo, MYQCOMMUNITY™, and the MYQCOMMUNITY logo are trademarks of Cytokinetics in the U.S.
© 2026 CYTOKINETICS. All Rights Reserved. US-MYQ-00312 05/2026

